




Images of the Month


Transition from summer to winter rainfall season

The month of April provided a wet cessation to a somewhat unpredictable 2023/24 summer rainfall season. Rainfall totals recorded over various provinces including North West, Gauteng, Mpumalanga, Free State, KwaZulu-Natal and the Eastern Cape, as well as the southern to eastern Western Cape, resulted in above-normal conditions, following relatively dry conditions during the latter part of the season. Wet conditions at the end of a growing season are typically not conducive to the ripening stages of the crop cycle and the commencement of harvest activities. However, greater parts of the summer rainfall region, including the western maize production area, can expect drier conditions during April-May-June as forecast by the South African Weather Service in March 2024 (<https://www.weathersa.co.za/home/seasonalclimate>).



Yet, in some areas, above-normal rainfall can be expected, potentially leading to crop damage or significant delays in harvest activities. Furthermore, winter conditions are typically characterized by cold and windy weather which could be devastating to livestock, especially during extreme weather events. Additionally, they may increase the likelihood of wildfires. Therefore, implementing preventative measures during this period is essential to mitigate these risks.

Expected Precipitation Conditions for AMJ 2024
Issued: Mar 2024



NATURAL RESOURCES AND ENGINEERING Soil, Climate and Water

CONTENTS:

- 1. Rainfall 2
- 2. Standardized Precipitation Index 4
- 3. Rainfall Deciles 6
- 4. Vegetation Conditions 7
- 5. Vegetation Condition Index 9
- 6. Vegetation Conditions & Rainfall 11
- 7. Fire Watch 15
- 8. Surface Water Resources 17
- 9. Agrometeorology 18
- 10. Geoinformatics 19
- 11. Analytical Laboratory 20
- 12. Microbiology Laboratory 21
- 13. CRID 22
- 14. Contact Details 22

Overview:

In April 2024, the country received notably higher rainfall compared to the preceding two months, particularly impacting the summer rainfall region. However, the Limpopo and Northern Cape provinces experienced below-normal rainfall during this period. The month began with warm temperatures, but by the 7th, an intense cut-off low brought heavy rainfall and strong winds to parts of the Western Cape, leading to destructive conditions, especially in the Overberg and southern Cape Winelands regions. Consequently, cold temperatures and rainy conditions prevailed over the interior and eastern regions of the country. On the 15th, severe thunderstorms caused disruptive flooding in the uGu and Ethekwini District Municipalities of KwaZulu-Natal.

Overall, significant rainfall was observed in the Highveld, southern interior and eastern parts of the Western Cape during April. Locations such as Vryburg, Ottosdal and Bloemfontein received up to 120 mm while Wepener, Kimberley and Lady Grey recorded over 150 mm for the month. Although the winter rainfall region also received above-average rainfall, the City of Cape Town experienced below-normal conditions.

1. Rainfall




Figure 1




Figure 2




Figure 3




Figure 4

Figure 1:

Widespread rainfall was experienced across South Africa during April 2024, with the highest amounts recorded over the interior, extending towards the Eastern and Western Cape provinces.

Figure 2:

Greater parts of the country received above-normal rainfall in April, while most of the Northern Cape, the City of Cape Town in the Western Cape, northern KwaZulu-Natal and Limpopo were mostly dry.

Figure 3:

Rainfall totals for the period between July 2023 and April 2024 indicate near- to above-normal rainfall over the country. Below-normal conditions can be observed in parts of the Northern Cape, North West, Limpopo, Free State and KZN.

Figure 4:

During February-April 2024, greater parts of the summer rainfall region and the southwestern parts of the Western Cape received significantly less rain compared to the same 3-month period last year. The rest of the country received relatively the same amount of rainfall, with isolated positive values noted in the Northern Cape.

Questions/Comments:

MasuphaE@arc.agric.za

Johan@arc.agric.za

2. Standardized Precipitation Index

Standardized Precipitation Index

The Standardized Precipitation Index (SPI - McKee *et al.*, 1993) was developed to monitor the occurrence of droughts from rainfall data. The index quantifies precipitation deficits on different time scales and therefore also drought severity. It provides an indication of rainfall conditions per quaternary catchment (in this case) based on the historical distribution of rainfall.

REFERENCE:

McKee TB, Doesken NJ and Kliest J (1993) The relationship of drought frequency and duration to time scales. In: Proceedings of the 8th Conference on Applied Climatology, 17-22 January, Anaheim, CA. American Meteorological Society: Boston, MA; 179-184.

The SPI maps revealing short-term (6-month), medium-term (12-month) and long-term (24- and 36-month) drought conditions ending in April 2024 are shown in Figures 5-8. The short-term SPI map indicates near-normal conditions across the country, with moderate drought notable over isolated parts of the summer rainfall region and the west coast. In contrast, the adjacent areas in Overberg, Cape Winelands and central Karoo experienced mild to severe wet conditions. The medium-term SPI map indicates moderate to extremely wet conditions over the Western Cape, isolated parts of the Eastern Cape, Northern Cape, Kwa-Zulu-Natal and Mpumalanga. The long-term SPI maps indicate predominantly wet conditions over greater parts of the country.

Questions/Comments:
MasuphaE@arc.agric.za
Johan@arc.agric.za




Figure 5




Figure 6




Figure 7




Figure 8

3. Rainfall Deciles

Deciles are used to express the ranking of rainfall for a specific period in terms of the historical time series. In the map, a value of 5 represents the median value for the time series. A value of 1 refers to the rainfall being as low or lower than experienced in the driest 10% of a particular month historically (even possibly the lowest on record for some areas), while a value of 10 represents rainfall as high as the value recorded only in the wettest 10% of the same period in the past (or even the highest on record). It therefore adds a measure of significance to the rainfall deviation.




Figure 9

Figure 9:

In April 2024, greater parts of the interior, moving towards the southwestern region of the country, experienced rainfall totals that compare well with historically wetter April months. The central parts of the Northern Cape, the City of Cape Town in the Western Cape, Limpopo, and northern parts of KwaZulu-Natal remained relatively dry.

Questions/Comments:
MasuphaE@arc.agric.za
Johan@arc.agric.za

Vegetation Mapping

The Normalized Difference Vegetation Index (NDVI) is computed from the equation:

$$NDVI = \frac{(IR - R)}{(IR + R)}$$

where:

IR = Infrared reflectance &
R = Red band

NDVI images describe the vegetation activity. A decadal NDVI image shows the highest possible "greenness" values that have been measured during a 10-day period.

Vegetated areas will generally yield high values because of their relatively high near infrared reflectance and low visible reflectance. For better interpretation and understanding of the NDVI images, a temporal image difference approach for change detection is used.

The Standardized Difference Vegetation Index (SDVI) is the standardized anomaly (according to the specific time of the year) of the NDVI.

4. Vegetation Conditions




Figure 10

Figure 10:

Compared to the historical averaged vegetation conditions, the 16-day NDVI map for April 2024 shows that the northeastern parts of central Limpopo, Free State and the Eastern Cape experienced below-normal vegetation activity, while the remaining parts of the country experienced above-normal conditions.

Figure 11:

The 16-day NDVI difference map for April 2024 compared to the preceding 16-day period shows that the western half of the country continued to experience normal to above-normal vegetation activity, while the eastern half experienced below-normal conditions.




Figure 11




Figure 12

Vegetation Mapping
(continued from p. 7)

Interpretation of map legend

NDVI-based values range between 0 and 1. These values are incorporated in the legend of the difference maps, ranging from -1 (lower vegetation activity) to 1 (higher vegetation activity) with 0 indicating normal/the same vegetation activity or no significant difference between the images.

Cumulative NDVI maps:

Two cumulative NDVI datasets have been created for drought monitoring purposes:

- Winter:** January to December
- Summer:** July to June




Figure 13

Figure 12: The 16-day NDVI difference map for April 2024 compared to the same period last year shows that the central and the northern parts of the country continued to experience poor vegetation conditions, while the western half experienced normal vegetation activity.

Figure 13: The Percentage of Average Seasonal Greenness (PASG) map for the past 4 months, compared to the long-term mean, shows that the country experienced a mixture of vegetation conditions with patches of both above-normal seasonal vegetation greenness and potential drought.

Questions/Comments:
MaakeR@arc.agric.za

5. Vegetation Condition Index

Vegetation Condition Index (VCI)

The VCI is an indicator of the vigour of the vegetation cover as a function of the NDVI minimum and maximum encountered for a specific pixel and for a specific period, calculated over many years.

The VCI normalizes the NDVI according to its changeability over many years and results in a consistent index for various land cover types. It is an effort to split the short-term weather-related signal from the long-term climatological signal as reflected by the vegetation. The VCI is a better indicator of water stress than the NDVI.

Vegetation Condition Index (VCI) for 29 Mar 2024 - 14 Apr 2024 compared to the long-term (22 years) mean




Figure 14

Figure 14:

The 16-day VCI map for April 2024 indicates that the western half of the Northern Cape experienced mainly below-normal vegetation conditions, while the eastern half experienced above-normal activity.

Figure 15:

The 16-day VCI map for April 2024 indicates that the western half of the Eastern Cape experienced above-normal vegetation conditions while the eastern half experienced mainly below-normal activity.

Vegetation Condition Index (VCI) for 29 Mar 2024 - 14 Apr 2024 compared to the long-term (22 years) mean




Figure 15




Figure 16




Figure 17

Figure 16:
The 16-day VCI map for April 2024 indicates that most parts of North West experienced below-normal vegetation conditions with patches of above-normal activity in isolated areas.

Figure 17:
The 16-day VCI map for April 2024 indicates that many parts of Gauteng continue to experience below-normal vegetation conditions, with patches of above-normal activity in the northern and central parts of the province.

Questions/Comments:
MaakeR@arc.agric.za

6. Vegetation Conditions & Rainfall

District Municipalities

- Towns
- District council
- Umkhanyakude
- Overberg
- iLembe
- King Cetshwayo
- Zululand
- Fezile Dabi
- Frances Baard
- Namakwa
- Pixley ka Seme
- ZF Mgcawu




Figure 18

Rainfall and NDVI Graphs

Figure 18: Orientation map showing the areas of interest for April 2024. The district colour matches the border of the corresponding graph.

Questions/Comments:
MaakeR@arc.agric.za

Figures 19-23: Indicate areas with higher cumulative vegetation activity for the last year.

Figures 24-28: Indicate areas with lower cumulative vegetation activity for the last year.




Figure 19




Figure 20




Figure 21







Figure 22






Figure 26




Figure 27




Figure 28

7. Fire Watch

Active Fires (Provided when data is available)

Forest and vegetation fires have temperatures in the range of 500 K (Kelvin) to 1000 K. According to Wien's Displacement Law, the peak emission of radiance for blackbody surfaces of such temperatures is at around 4 μm . For an ambient temperature of 290 K, the peak of radiance emission is located at approximately 11 μm . Active fire detection algorithms from remote sensing use this behaviour to detect "hot spot" fires.

Figure 29:

The graph shows the total number of active fires detected from 29 March to 30 April 2024 per province. Fire activity was higher in Mpumalanga, Limpopo and the Western Cape compared to the long-term average.




Figure 29

Active fires detected between 29 March - 30 April 2024




Figure 30

Figure 30:

The map shows the location of active fires detected between 29 March and 30 April 2024.




Figure 31: The graph shows the total number of active fires detected from 1 January to 30 April 2024 per province. Fire activity was higher in the Eastern Cape, Free State and North West compared to the long-term average.

Figure 31




Figure 32: The map shows the location of active fires detected between 1 January and 30 April 2024.

Questions/Comments:
MaakeR@arc.agric.za

Figure 32

Countrywide surface water areas (SWAs) are mapped on a monthly basis by GeoTerraImage using Sentinel-2 satellite imagery from the start of its availability at the end of 2015.

Figure 33 represents a comparison between the area of water available now and the maximum area of surface water recorded in the last 8 years. This 8-year historical window represents the operational period of the satellite from which the water information has been generated. In this map, any value less than 100 represents water catchments within which the current month's total surface water is less than the maximum extent recorded for the same area since the end of 2015.



Figure 34 represents a comparison between the area of surface water now and for the same month last year. In this map, any value less than 100 represents water catchments within which the current month's total surface water is less than that recorded in the same water catchment, in the same month, last year.

The long-term map for April 2024 shows a marked increase, countrywide, of the number of catchments now representing 80-100% levels, indicative of better April rainfall conditions this year, compared to previous years.

The comparison between April 2024 and April 2023 is now indicating improvements in water conditions compared to last year in several Karoo and east coast catchments (150% to 200 plus % levels). The remainder of the country shows little change compared to the conditions last year (50-150% levels).

The SWA maps are derived from the monthly data generated and available through GeoTerraImage's 'Msanzi Amanzi' web information service: <https://www.water-southafrica.co.za>

Questions/Comments:
mark.thompson@geoterraimage.com





Agrometeorology

The programme uses weather and climate information for agricultural planning and the enhancement of crop and livestock production systems. The impact of climate variability and change in the agricultural sector is investigated. Due to the increasing pressure to reduce greenhouse gas emissions globally, climate change mitigation is also an important facet of our activities. The Weather Station Network and Climate Database are maintained as a national asset for the benefit of the agriculture sector.

Activities

Agrometeorology and Crop Modelling

- Assessing climate risk for an area in relation to a particular crop
- Agroclimatological analysis of the suitability for crop production at a particular location
- Development of early warning systems for climate hazards (e.g. drought, floods)
- Agrometeorological forecasting and advisory services
- Crop modelling to assess the impact of weather conditions and climate on agriculture
- Conducting crop yield forecasting exercises, hydrological modelling, hydrometeorology and biometereology studies

Climate Change Adaptation and Mitigation

- Conducting research on possible impact of projected climate change on agricultural activities, potential, greenhouse gas emissions from various land use, climate change, mitigation and adaptation strategies for agriculture
- Developing greenhouse gas inventories at farm and national levels
- Conducting research on climate change mitigation and adaptation strategies for agriculture
- Promoting low-carbon technologies

Climate Monitoring, Products and Services

- Developing and maintaining a network of over 500 weather stations distributed all over the country
- Archiving historical and current weather data of good quality with some datasets dating back to 1900
- Developing weather/climate products and services together with stakeholders and clients to meet their specific requirements
- Disseminating weather/climate data, products and services via multiple platforms

Contact Person: Dr Mokhele Moeletsi

Tel: 012 310 2537 • E-mail: moeletsim@arc.agric.za

ARC-Natural Resources and Engineering
Soil, Climate and Water Campus
 600 Belvedere Street, Arcadia, 0083
 Private Bag X79, Pretoria 0001
 Tel: 012 310 2500 • Fax: 012 323 1157
 Website: www.arc.agric.za



For more information contact:
 Adri Laas - Public Relations Officer
 E-mail: adril@arc.agric.za

SOIL, CLIMATE AND WATER



GeoInformatics

The programme focuses on applied Geographical Information Systems (GIS) and provides leadership in GIS products, solutions and decision support systems for agriculture and natural resources management. The Coarse Resolution Satellite Image Archive and Information Database is maintained as a national asset.

Activities

Digital/Smart Agriculture/Drone Platform - Applications

- Yield & production estimation
- Insurance index
- Mapping crop types
- Monitoring growth stages
- Weed/invasive sp. mapping
- Water requirement
- Smart & digital agriculture
- Disease/pests



Applications in Natural Resources/National Assets

- Early warnings
- National & Provincial advisories
- Crop suitability changes
- Crop statistics
- Crop stress
- Spatially explicit information dissemination systems, e.g. Umlindi newsletter



Applications in Rangelands, Livestock and Wildlife

- Early warnings
- National & Provincial advisories
- Rangeland suitability
- Rangeland dynamics
- Rangeland stresses
- Spatially explicit information dissemination systems, e.g. Umlindi newsletter



Contact Person: Dr George Chirima
Tel: 012 310 2672 • E-mail: chirimaj@arc.agric.za

ARC-Natural Resources and Engineering
Soil, Climate and Water Campus
600 Belvedere Street, Arcadia, 0083
Private Bag X79, Pretoria 0001
Tel: 012 310 2500 • Fax: 012 323 1157
Website: www.arc.agric.za



For more information contact:
Adri Laas - Public Relations Officer
E-mail: adri@arc.agric.za



SOIL, CLIMATE AND WATER



Analytical Laboratory

The unit focuses on the various procedures to analyze and determine the properties of soil, water and associated materials, mainly for agricultural purposes. The laboratory operates a range of equipment and participates in various quality control schemes, both local and international. The water analysis for anions is SANAS-accredited and other accreditations are underway.

Analyses and Services

Soil Physical Analysis

- Texture (sand, silt and clay content)
- Water-holding capacity
- Soil moisture content
- Bulk density
- Shrink-swell capacity

Soil Chemical Analysis

- pH
- Exchangeable and extractable cations
- Acidity
- Soil Organic Carbon
- Nitrogen content and C/N ratio
- Phosphorus
- Micronutrients

Soil Fertility

- Analysis package for farmers & gardeners
- Fertilizer recommendations for specific crops

Water Analysis

- pH, EC, anions, cations
- Water quality

ICP Scan

- Semi-quantitative scan for a range of elements (Li, Be, Ti, V, Cr, Mn, Co, Ni, Cu, Zn, As, Se, Rb, Sr, Mo, Cd, Sn, Sb, Te, Cs, Ba, La, W, Pt, Hg, Tl, Pb, Bi, U), can be done on soil, water and plant

Plant Material Analysis

For example: leaves, roots, growth media, etc. – drying, milling, pH, EC, C, N, nutrients and toxic elements

Special Sample Analysis

- For example: sludges, compost, fertilizers – composition and other properties
- Elemental analysis of animal tissue (e.g. hair, bones, liver, muscle, milk)

For more information or to obtain prices or quotation, contact the Laboratory Manager: Ms. Zanele Hlam
Tel: 012 310 2531 • E-mail: HlamZ@arc.agric.za

In order to assist clients who wish to send samples to ARC, the courier costs can be borne by ARC for analysis packages of R10 000 or more.

Contact the Laboratory Manager for details.



**ARC-Natural Resources and Engineering
Soil, Climate and Water Campus**

600 Belvedere Street, Arcadia, 0083 • Private Bag X79, Pretoria 0001

Tel: 012 310 2500 • Fax: 012 323 1157

Website: www.arc.agric.za

For more information contact:

Adri Laas - Public Relations Officer • E-mail: adri@arc.agric.za

SOIL, CLIMATE AND WATER



Microbiology and Environmental Biotechnology Laboratory

The Microbiology and Environmental Biotechnology Research Group forms part of the Soil Science Programme at ARC-SCW. The research group utilizes both fundamental as well as applied microbiology and biotechnology approaches to address soil, climate and water related problems in a sustainable and eco-friendly manner.

Analyses and Services

Renewable energy generation

- Gas Chromatography analysis of biogas - methane and carbon dioxide content measurements

Nanotechnology

- UV-Visible spectrophotometer analysis for colloidal nanoparticle synthesis

Phytochemical extraction

- Hotplate extraction of phytochemicals
- Soxhlet extraction of phytochemicals
- Microwave-assisted extraction of phytochemicals

Community-Level Physiological Profiling (CLPP)

- Microbial functional analysis using Biolog 31C plates

For information on microbiological analyses contact

Dr Ashira Roopnarain

Tel: 012 310 2650 • E-mail: RoopnarainA@arc.agric.za

In order to assist clients who wish to send samples to ARC, the courier costs can be borne by ARC for analysis packages of R10 000 or more.

Contact the Laboratory Manager for details.

**ARC-Natural Resources and Engineering
Soil, Climate and Water Campus**

600 Belvedere Street, Arcadia, 0083 • Private Bag X79, Pretoria 0001

Tel: 012 310 2500 • Fax: 012 323 1157

Website: www.arc.agric.za

For more information contact:

Adri Laas - Public Relations Officer • E-mail: adri@arc.agric.za



The Coarse Resolution Imagery Database (CRID)

NOAA AVHRR

ARC-NRE has an archive of daily NOAA AVHRR data dating from 1985 to 2004. This database includes all 5 bands as well as the Normalized Difference Vegetation Index (NDVI), Active Fire and Land Surface Temperature (LST) images. The NOAA data are used, for example, for crop production and grazing capacity estimation.

MODIS

MODIS data is distributed by the Land Processes Distributed Active Archive Center (LP DAAC), located at the U.S. Geological Survey's EROS Data Center. The MODIS sensor is more advanced than NOAA with regard to its high spatial (250 m² to 1 km²) and spectral resolution. ARC-NRE has an archive of MODIS (version 4 and 5) data.

- MODIS v4 from 2000 to 2006
- MODIS v5 from 2000 to present

Datasets include:

- MOD09 (Surface Reflectance)
- MOD11 (Land Surface Temperature)
- MOD13 (Vegetation Products)
- MOD14 (Active Fire)
- MOD15 (Leaf Area Index & Fraction of Photosynthetically Active Radiation)
- MOD17 (Gross Primary Productivity)
- MCD43 (Albedo & Nadir Reflectance)
- MCD45 (Burn Scar)

Coverage for version 5 includes South Africa, Namibia, Botswana, Zimbabwe and Mozambique.

More information:

<http://modis.gsfc.nasa.gov>

VG4AFRICA and GEOSUCCESS

SPOT NDVI data is provided courtesy of the VEGETATION Programme and the VGT4AFRICA project. The European Commission jointly developed the VEGETATION Programme. The VGT4AFRICA project disseminates VEGETATION products in Africa through GEONETCast.

ARC-NRE has an archive of VEGETATION data dating from 1998 to the present. Other products distributed through VGT4AFRICA and GEOSUCCESS include Net Primary Productivity, Normalized Difference Wetness Index and Dry Matter Productivity data.

Meteosat Second Generation (MSG)

ARC-NRE has an operational MSG receiving station. Data from April 2005 to the present have been archived. MSG produces data with a 15-minute temporal resolution for the entire African continent. Over South Africa the spatial resolution of the data is in the order of 3 km. ARC-NRE investigated the potential for the development of products for application in agriculture. NDVI, LST and cloud cover products were some of the initial products derived from the MSG SEVIRI data. Other products derived from MSG used weather station data, including air temperature, humidity and solar radiation.

Rainfall maps

- Combined inputs from 450 automatic weather stations from the ARC-NRE Soil, Climate and Water weather station network, 270 automatic rainfall recording stations from the South African Weather Service (SAWS), satellite rainfall estimates from the Famine Early Warning System Network: <http://earlywarning.usgs.gov> and long-term average climate surfaces developed at the ARC-NRE.

Solar Radiation and Evapotranspiration maps

- Combined inputs from 450 automatic weather stations from the ARC-NRE Soil, Climate and Water weather station network.
- Data from the METEOSAT Second Generation (MSG) 3 satellite via GEONETCAST: <http://www.eumetsat.int/website/home/Data/DataDelivery/EUMETCast/GEONETCast/index.html>.



NATURAL RESOURCES AND ENGINEERING Soil, Climate and Water

Private Bag X79, Pretoria 0001,
South Africa
600 Belvedere Street, Arcadia, Pretoria, South Africa

Reneilwe Maake

Project Leader: Coarse Resolution Imagery
Database (CRID)
Tel: 012 310 2533
E-mail: MaakeR@arc.agric.za

The operational Coarse Resolution Imagery Database (CRID) project of ARC-NRE is funded by the Department of Agriculture, Land Reform and Rural Development (DALRRD). Development of the monitoring system was made possible at its inception through LEAD funding from the Department of Science and Technology.

For further information please contact:
Reneilwe Maake – 012 310 2533, MaakeR@arc.agric.za
Adri Laas – 012 310 2518, AdriL@arc.agric.za

To subscribe to the newsletter, please submit a request to:
MaakeR@arc.agric.za

What does Umlindi mean?
UMLINDI is the Zulu word for “the watchman”.

DISCLAIMER:

The ARC-NRE and its collaborators have obtained data from sources believed to be reliable and have made every reasonable effort to ensure accuracy of the data. The ARC-NRE and its collaborators cannot assume responsibility for errors and omissions in the data nor in the documentation accompanying them. The ARC-NRE and its collaborators will not be held responsible for any consequence from the use or misuse of the data by any organization or individual.